

HOW TO USE

INVELTYS® (loteprednol etabonate ophthalmic suspension) 1%

**For complete instructions on how to use INVELTYS,
please see the Full Prescribing Information.**

1. If you wear contact lenses, remove them before using INVELTYS.



2. There are two caps on your bottle of INVELTYS. Hold the bottle firmly by its neck. Remove the WHITE CAP by twisting it clockwise. You can throw this cap away.

3. Shake the bottle for 1 to 2 seconds.

4. Then, remove the PINK CAP by turning it counter-clockwise. Keep this cap. DO NOT ALLOW THE TOP OF THE DROPPER TO TOUCH ANY SURFACE.



5. Hold the bottle directly above your eye. Squeeze the middle of the bottle gently to put 1 to 2 drops (follow your doctor's recommendation) into your eye.



6. Place the pink cap back onto the bottle and tighten.

7. If you use contact lenses, wait for 15 minutes before placing them back in.

What is INVELTYS?

INVELTYS (loteprednol etabonate ophthalmic suspension) 1% is a prescription eyedrop medication used to treat inflammation and pain after ocular surgery.

Important Safety Information

Who should NOT use INVELTYS?

- You should NOT use INVELTYS if you have an infection in your eye.

Please see additional Important Safety Information on reverse.

Please see accompanying Full Prescribing Information.

INVELTYS®
(loteprednol etabonate
ophthalmic suspension) 1%

Important Safety Information (continued)

What should I tell my eye care professional before taking INVELTYS?

- Before taking INVELTYS, tell your doctor about all your medical conditions, including if you have an eye infection or a history of herpes infections.

What are the possible side effects of INVELTYS?

- Prolonged use of corticosteroids, such as INVELTYS, may result in high eye pressure (glaucoma) with damage to the optic nerve, affecting your vision. If you use INVELTYS for longer than 10 days, your doctor may monitor pressure in your eye.
- Use of corticosteroids may result in cataract formation and may delay healing after cataract surgery.
- Prolonged use of corticosteroids may increase the risk of eye infections.

What are the most common side effects of INVELTYS?

- In clinical trials, the most common side effects were eye pain and clouding of the lens capsule (posterior capsular opacification). These effects may have been due to the surgery.

How should I take INVELTYS?

- Take INVELTYS as directed by your doctor.
- Contact lenses should be removed prior to instillation of INVELTYS and may be reinserted 15 minutes following administration. The preservative in INVELTYS may be absorbed by soft contact lenses.
- Shake the bottle for one to two seconds before using INVELTYS.
- Instill one to two drops of INVELTYS into the affected eye twice daily beginning a day after surgery throughout the first 2 weeks after surgery, or as directed.
- Do not allow the dropper tip to touch any surface, as this may contaminate the eye drop.
- If pain develops, or if redness, itching, or inflammation becomes worse, you should consult your doctor.

Please see accompanying Full Prescribing Information.



INVELTYS is a registered trademark of Kala Pharmaceuticals.

©2019 Kala Pharmaceuticals

Printed in the USA. US-INV-1900109 May 2019 www.inveltys.com

INVELTYS®
(loteprednol etabonate
ophthalmic suspension) 1%